



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

tude  $3^m.0$ , thus yielding a mean value of  $0''.0100$  for the parallax of an eighth magnitude star. According to Kapteyn's formula  $\bar{\pi} = 0''.0158 (0.78)^{m-5.5}$  the mean parallax for a star of visual magnitude  $8^m.0$  is  $0''.0085$ . Although the agreement is very close, we must not overestimate its weight, on account of the uncertainty of the color-index for the stars with high luminosity. If, for instance, we disregard the first three dots in figure 2, we can, without straining the figures, arrive at a color-index of  $0^m.56$  for  $M = 3^m.8$ , corresponding to a parallax of  $0''.0144$  or 70% more than Kapteyn's value. A color-index of  $0^m.28$  (F0) might be reached at  $m = 1^m.0$  and  $M = 0^m.0$  giving a mean parallax of  $0''.063$  for a first magnitude star, whereas Kapteyn finds  $0''.048$ .

1922, January 3.

WILLEM J. LUYTEN.

---

#### REMARKS ON H. D. CURTIS' PAPER ON STELLAR LUMINOSITIES

The interesting paper by Heber D. Curtis<sup>1</sup> in the last number of this journal, with its excellent material of absolute magnitudes, suggests a few comments. First, it may be pointed out that the well-known relative scarcity of F and early G giants is even more striking when compared with the excess of "super-giants" of these classes (spectroscopic absolute magnitudes for *Cepheids*). This tends to make the diagram look as if the distribution of absolute magnitudes follows two parallel lines. The higher one of these two lines may link up with the B-supergiants, such as  $\beta$  *Orionis*,  $\alpha$  *Crucis*, and the Walf-Rayet stars, to some of which Lundmark<sup>2</sup> assigns absolute magnitudes of  $-4^m.0$ .

In connection with the probable parallaxes of a  $9^m.5$  star, which Curtis derives on page 38 of his paper, it may be remarked that, although in space the dwarfs greatly outnumber the giants, this may not be applied when dealing with stars of the same apparent magnitude. If we take the number of giants to be only 1% of the total, and assume the mean absolute magnitudes of giants and dwarfs to be  $+0^m.8$  and  $+6^m.0$  respectively (which holds for the K-stars), both with a Gaussian distribution, then

<sup>1</sup>Publ. A. S. P. **34**, 33, 1922.

<sup>2</sup>Publ. A. S. P. **34**, 40, 1922.

the dwarfs form only 40% of the total number at apparent magnitude  $9^m.9$ , and 18% at  $6^m.0$ .

WILLEM J. LUYTEN.

#### ADDITIONAL NOTE ON FAINT EARLY-TYPE STARS WITH LARGE PROPER-MOTIONS

Besides the list already published in this journal,<sup>1</sup> three even more striking cases have been found of white stars with low apparent magnitude and large proper-motion.

| NAME                               | R.A.          | 1900.0 DEC.      | MAG.     | SP. | P.M.     |
|------------------------------------|---------------|------------------|----------|-----|----------|
| Wolf 1056 . . . . .                | $0^h33^m29^s$ | $+29^\circ47'.3$ | $11^m.0$ | A3  | $1''.69$ |
| C. P. D.— $47^\circ7499$ . . . . . | 15 50 15      | $-47^\circ20.6$  | 9 .5     | B9  | 0 .85    |
| C. P. D.— $40^\circ7356$ . . . . . | 16 25 32      | $-40^\circ6.0$   | 10 .1    | A0  | 1 .05    |

The spectrum of the first star was determined by Wolf,<sup>2</sup> but his estimate may differ materially from the Harvard classification. For the two C. P. D. stars, no stars of the same brightness are close to them. If we assume that these stars have the mean absolute magnitude of a class A0 star, *i. e.*,  $+1^m.2$ , then the linear motions in space corresponding to the proper-motions are 7300, 1800, 3000 km. per sec. respectively. If on the other hand, we assume the velocity not to be higher than 100 km. per sec., which is still very high for A-stars, we find as upper limits for the absolute magnitudes  $+10^m.5$ ,  $+7^m.5$ , and  $+8^m.6$  respectively, quite comparable with that of  $\alpha^2$  Eridani B ( $+11^m.2$ ).

These considerations make it probable that the three early-type stars are dwarfs, and also that they will prove interesting objects for both radial velocity and parallax observers in the southern hemisphere.

1922, March 3.

WILLEM J. LUYTEN.

#### THE TOTAL RADIATION FROM $\alpha$ CETI

In the course of some stellar observations with vacuum thermocouples, measures have been made with the Hooker telescope of the heat received from  $\alpha$  Ceti. On December 6, 1921, the

<sup>1</sup>Publ. A. S. P., **34**, 54, 1922.

<sup>2</sup>A. N., **210**, 293, 1919.